

Vasco VIEIRA SERRÃO  
Andréa MARTINS  
M. João Paiva LOPES

Dermatology Department,  
Hospital dos Capuchos, Alameda Santo  
António dos Capuchos,  
1150 – 314 Lisboa, Portugal

**Reprints:** V. Vieira Serrão  
<vvvsferreira@netcabo.pt>

## Infliximab in recalcitrant generalized pustular arthropathic psoriasis

Generalized pustular psoriasis is an unstable inflammatory type of psoriasis, with widespread areas of erythema and sterile pustules, associated with fever and systemic symptoms. Infliximab is a monoclonal antibody with anti-TNF $\alpha$  activity, approved for use in psoriasis. We describe a male patient with a long history of stable arthropathic psoriasis, hospitalized with a generalized pustular psoriasis and acute exacerbation of articular complaints. The disease was resistant to multiple therapies (acitretin, methotrexate and corticosteroids), so the patient was started on infliximab, with a very rapid response of both cutaneous and articular symptoms. He had complete clearing of lesions at week 12, and marked improvement of the articular symptoms. No recurrence occurred at 8 months of follow-up with infliximab every 8 weeks. Infliximab had an extremely rapid therapeutic action response on a recalcitrant generalized pustular psoriasis. The articular response was also excellent, with significant improvement of quality of life.

**Key words:** anti-TNF-alpha, arthropathic, biologics, infliximab, pustular psoriasis

Article accepted on 22/9/2007

**G**eneralized pustular psoriasis (Von Zumbusch type) is an unstable inflammatory type of psoriasis, with widespread areas of erythema and sterile pustules, associated with fever and systemic symptoms [1]. In this form of psoriasis, infiltration of the skin with neutrophils plays an important role, leading to the characteristic sterile pustules [2]. Infliximab is a monoclonal antibody with activity against tumour necrosis factor- $\alpha$  (TNF- $\alpha$ ), approved for use in psoriasis [3]. The very rapid onset of action and reported efficacy makes it an appealing option for the treatment of generalized pustular psoriasis [4].

### Case report

A 59-year-old male patient, with a 24-year history of plaque psoriasis with moderate articular involvement, was hospitalized with a severe generalized pustular eruption. The disease had been stable and localized, with moderate articular complaints, mostly axial. Betamethasone cream, calcipotriol ointment and naproxen were the only previous medication, along with chronic anti-hypertensive therapy (indapamide, losartan and perindopril).

He developed an inflammatory flare-up of the disease, with erythematous plaques on the upper trunk, which rapidly generalized, affecting 90% of the body surface (*figure 1*). Small non-follicular pustules developed over these inflammatory lesions and fever (38 °C), malaise, and severe worsening of articular complaints occurred simultaneously, causing notable difficulty with mobilization.

The patient denied previous use of systemic corticosteroids or any new medication recently. Infectious and metabolic

triggers were ruled out by clinical and analytical evaluation, in particular hypocalcemia, streptococcal or human immunodeficiency virus infection. On admission, the patient had a white blood cell count of  $13.4 \times 10^3/\mu\text{L}$  with 91% neutrophils, erythrocyte sedimentation rate of 58 mm and C reactive protein of 23.1 mg/dL.

Treatment was started with methotrexate (15 mg/weekly PO) and acitretin (35 mg/d), but new pustules and very severe articular pain continued after three weeks of treatment. As uncontrolled hypertension precluded the use of cyclosporine, systemic corticosteroids (1 mg/kg/d) were added, while waiting for the full therapeutic effect of methotrexate and acitretin. There was a rapid resolution of the pustular eruption, fever and improvement of articular complaints, and slow tapering of corticosteroids was started (10% weekly dose reduction over 5 weeks). When the dose reached 0.5 mg/kg/d, new pustules appeared together with worsening of articular complaints. The corticosteroids were stopped, and after exclusion of latent tuberculous infection, infliximab was started at 5 mg/kg IV (weeks 0, 2 and 6, and then every 8 weeks) and both methotrexate and acitretin were maintained. There was an extremely rapid response, noticeable from the second day, with complete clearance of the pustular eruption at the end of the first week (*figure 2*). There was also an improvement in both articular pain and mobility, and methotrexate and acitretin were stopped at the second infliximab infusion. Complete clearance of cutaneous lesions was achieved at 12 weeks, with no significant side effects reported. No relapse was noted at 26 weeks follow-up, with special emphasis on the continued articular improvement.



**Figure 1.** Generalized pustular psoriasis. **A)** Multiple active lesions despite combination therapy; **B)** Rapid improvement after one week of the first infliximab infusion; **C)** Complete resolution after 3 infliximab infusions (12 weeks).

## Discussion

Generalized pustular psoriasis (GPP) is a serious dermatological disease, which can result in significant morbidity and even mortality [5]. The disease is characterized by fever, systemic symptoms and generalized pustule formation on the skin [1].

GPP frequently occurs as an inflammatory flare-up of a stable psoriasis, often associated with the use of systemic corticosteroids, and more rarely topical corticosteroids [6, 7]. Many other triggers have been described, like pregnancy, hypocalcemia, infections and sudden withdrawal of anti-psoriatic therapy [8-10]. No infectious, metabolic or drug-induced trigger was identified in our patient.

Acitretin has good efficacy in pustular psoriasis, with inhibition of neutrophil migration and activation [11]. We com-

bined acitretin and methotrexate for our patient in order to target both the skin and articular involvement [12]. Systemic corticosteroids were used to rapidly control the inflammatory flare of the disease, while waiting for the full therapeutic effect of methotrexate.

Infliximab has shown good efficacy and very rapid onset of action for plaque psoriasis and psoriatic arthritis, with a beneficial impact on quality of life and work productivity [3, 13-15]. The experience with infliximab in clinical practice has been expanding, although few reports have been published on its use in GPP [16-19]. Even in severe recalcitrant cases, infliximab has been shown to be highly effective in rapidly controlling the disease [5, 20-22]. By inhibiting TNF- $\alpha$ , it appears to downregulate neutrophil-attractant chemokines which play a relevant role on the pathogenesis of the disease, such as interleukin-8 (IL-8),



**Figure 2.** Generalized pustular psoriasis and infliximab therapy: **A)** Pre-treatment; **B)** After one week; **C)** After 12 weeks.

growth-related oncogene (Gro- $\alpha$ ) and monocyte chemoattractant protein (MCP-1) [22].

Our patient had an excellent response to infliximab, with improvement noticeable from the second day. Complete clearance of the lesions occurred at week 12, after only three infusions. This very rapid onset of action has already been described [20, 21], making it a very attractive therapy for inpatients, by rapidly and efficiently controlling the disease and allowing early discharge from hospital [4]. ■

**Acknowledgements.** Financial support: None. Conflict of interest: None.

## References

1. Griffiths CE, Christophers E, Barker JN, Chalmers RJ, Chimenti S, Krueger GG, Leonardi C, Menter A, Ortonne JP, Fry L. A classification of psoriasis vulgaris according to phenotype. *Br J Dermatol* 2007; 156: 258-62.
2. Iizuka H, Takahashi H, Ishida-Yamamoto A. Pathophysiology of generalized pustular psoriasis. *Arch Dermatol Res* 2003; 295(Suppl 1): S55-S59.
3. Reich K, Nestle FO, Papp K, Ortonne JP, Evans R, Guzzo C, Li S, Dooley LT, Griffiths CE. EXPRESS study investigators. Infliximab induction and maintenance therapy for moderate-to-severe psoriasis: a phase III, multicentre, double-blind trial. *Lancet* 2005; 366: 1367-74.
4. Nelson AA, Pearce DJ, Fleischer Jr. AB, Balkrishnan R, Feldman SR. Inpatient management of severe psoriasis. *J Drugs Dermatol* 2005; 4: 564-70.
5. Lewis TG, Tuchinda C, Lim HW, Wong HK. Life-threatening pustular and erythrodermic psoriasis responding to infliximab. *J Drugs Dermatol* 2006; 5: 546-8.
6. Elston GE, Charles-Holmes R, Carr RA. Precipitation of generalized pustular psoriasis by prednisolone. *Clin Exp Dermatol* 2006; 31: 133-4.
7. Telfer NR, Dawber RP. Generalized pustular psoriasis associated with withdrawal of topical clobetasol-17-propionate. *J Am Acad Dermatol* 1987; 17: 144-5.
8. Hong SB, Kim NI. Generalized pustular psoriasis following withdrawal of short-term cyclosporin therapy for psoriatic arthritis. *J Eur Acad Dermatol Venereol* 2005; 19: 522-3.
9. Cassandra M, Conte E, Cortez B. Childhood pustular psoriasis elicited by the streptococcal antigen: a case report and review of the literature. *Pediatr Dermatol* 2003; 20: 506-10.
10. Gaylor ML, Duvic M. Generalized pustular psoriasis following withdrawal of efalizumab. *J Drugs Dermatol* 2004; 3: 77-9.
11. Lee CS, Koo J. A review of acitretin, a systemic retinoid for the treatment of psoriasis. *Expert Opin Pharmacother* 2005; 6: 1725-34.
12. Lebwohl M, Menter A, Koo J, Feldman SR. Combination therapy to treat moderate to severe psoriasis. *J Am Acad Dermatol* 2004; 50: 416-30.
13. Rinaldi F, Provenzano G, Termini A, Spinello M, La Seta F. Long term infliximab treatment for severe psoriatic arthritis: evidence of sustained clinical and radiographic response. *Ann Rheum Dis* 2005; 64: 1375-6.
14. Katugampola RP, Lewis VJ, Finlay AY. The Dermatology Life Quality Index: assessing the efficacy of biological therapies for psoriasis. *Br J Dermatol* 2007; 156: 945-50.
15. Reich K, Nestle FO, Wu Y, Bala M, Eisenberg D, Guzzo C, Li S, Dooley LT, Griffiths CE. Infliximab treatment improves productivity among patients with moderate-to-severe psoriasis. *Eur J Dermatol* 2007; 17: 381-6.
16. Ozawa A, Ohkido M, Haruki Y, Kobayashi H, Ohkawara A, Ohno Y, Inaba Y, Ogawa H. Treatments of generalized pustular psoriasis: a multicenter study in Japan. *J Dermatol* 1999; 26: 141-9.
17. Augéy F, Renaudier P, Nicolas JF. Generalized pustular psoriasis (Zumbusch): a French epidemiological survey. *Eur J Dermatol* 2006; 16: 669-73.
18. Roelofzen JH, Aben KK, Khawar AJ, Van de Kerkhof PC, Kiemeneij LA, Van Der Valk PG. Treatment policy for psoriasis and eczema: a survey among dermatologists in the Netherlands and Belgian Flanders. *Eur J Dermatol* 2007; 17: 416-21.
19. Trent JT, Kerdell FA. Successful treatment of Von Zumbusch pustular psoriasis with infliximab. *J Cutan Med Surg* 2004; 8: 224-8.
20. Lisby S, Gniadecki R. Infliximab (Remicade) for acute, severe pustular and erythrodermic psoriasis. *Acta Derm Venereol* 2004; 84: 247-8.
21. Newland MR, Weinstein A, Kerdell F. Rapid response to infliximab in severe pustular psoriasis, von Zumbusch type. *Int J Dermatol* 2002; 41: 449-52.
22. Benoit S, Toksoy A, Brocker EB, Gillitzer R, Goebeler M. Treatment of recalcitrant pustular psoriasis with infliximab: effective reduction of chemokine expression. *Br J Dermatol* 2004; 150: 1009-12.